

How Sensitive are Tail-Related Risk Measures in a Contamination Neighbourhood?

Wolfgang Karl Härdle
Chengxiu Ling

Ladislaus von Bortkiewicz Chair of Statistics
Humboldt–Universität zu Berlin
School of Mathematics and Statistics
Southwest University
<http://lwb.wiwi.hu-berlin.de>
<http://swu.edu.cn>



Motivation

- Financial asset returns often contain observations that are inconsistent with the majority of the data
- Estimation or mis-specification errors in the portfolio loss distribution can have a considerable impact on risk measures
- Choosing risk measures and its risk level plays an important role in financial practice and risk management

Example

CRIX, the benchmark cryptocurrency index by Trimborn and Härdle (2016)




Figure 1: Time series (left) and normal Q-Q plot (right) of the daily log returns of CRIX during 20140731-20180101, standardized by using GARCH(1, 1) model. Data source: crix.berlin

Common risk measures

- ▣ **Value-at-Risk (VaR)**, the minimal loss of the α worst cases
- ▣ **Expected-shortfall (ES)**, the expectation of the α worst loss
- ▣ **Expectile**, one-to-one mapping with VaR, reflects the tail heaviness through the **expectile-quantile transformation**

- ▶ Expectile level w_α such that $e_{w_\alpha} = q_\alpha$, satisfies [Details](#)

$$w_\alpha = \frac{\int_{-\infty}^u x dF(x) - u\alpha}{2\{\int_{-\infty}^u x dF(x) - u\alpha\} + u - E[X]}, \quad u = q_\alpha \quad (1)$$

- ▶ ES using expectile, Taylor (2008)

$$ES_\alpha = e_{w_\alpha} + \frac{e_{w_\alpha} - E[X]}{1 - 2w_\alpha} \frac{w_\alpha}{\alpha}$$

Example

Consider a mixture model given in Kuan et al. (2009)

$$F_{\epsilon}(x) = (1 - \epsilon)\Phi(x\sqrt{1 - \epsilon}) + \epsilon\Phi(x\sqrt{\epsilon})$$

- Contamination level ϵ
- Heaviness parameter $\sigma = 1/\sqrt{\epsilon}$
- Interplay of risk level α and contamination level ϵ to determine risk measures

Research Questions

- How to establish suitable contamination models for such data?
- How sensitive are risk measures for such models?
- How are risk measures adjusted for contamination data?

Outline

1. Motivation ✓
2. Contamination models
3. Approximation results
4. Numerical examples
5. Empirical study
6. Conclusions

Contamination Models

Consider

$$F_\epsilon(x) = (1 - \epsilon)F(x) + \epsilon H(x), \quad x \in \mathbb{R}, \quad \epsilon \in [0, 1]$$

- F stands for the pre-supposed ideal model
- H represents plausible deviations from F
- ϵ reflects the amount of uncertainty in F

Common Assumptions

- Both F and H have infinite left endpoints since we are interested with infinite risks
- Catastrophe contamination attracts financial regulators and thus assume H has a heavier tail than F
- Risk measures concern extreme value risk management and thus suppose F and H are common distributions in EVT

Notation

- Recall that F and F_ϵ stand for the pre-supposed ideal model and the contamination model in the ϵ neighbourhood of F
- Denote by ϱ_α and $\varrho_\alpha(\epsilon)$ the risk measure ϱ of F and F_ϵ
 - ▶ Value-at-Risk (VaR): q_α and $q_\alpha(\epsilon)$
 - ▶ Expected shortfall (ES): ES_α and $ES_\alpha(\epsilon)$
 - ▶ Expectile-quantile transformation level: w_α and $w_\alpha(\epsilon)$

Approximation of Risk Measures

- **Case A:** common contamination with $\epsilon \in (0, 1]$ fixed and $\alpha \rightarrow 0$
- **Case B:** interplay between ϵ and α with $\epsilon = \epsilon_\alpha \rightarrow 0$ as $\alpha \rightarrow 0$
- **Case C:** infinitesimal contamination with $\alpha \in (0, 1)$ fixed and $\epsilon \rightarrow 0$

Case A: Fixed ϵ

Theorem A Let $F_\epsilon(x) = (1 - \epsilon)F(x) + \epsilon H(x)$ with $\epsilon \in (0, 1]$. We have as $\alpha \rightarrow 0$

$$q_\alpha(\epsilon) \sim q_{\alpha'}(1) \quad \text{with} \quad \alpha' \stackrel{\text{def}}{=} \alpha/\epsilon$$

Further, if both exist with finite means, then

$$ES_\alpha(\epsilon) \sim ES_{\alpha'}(1), \quad \frac{w_\alpha(\epsilon)}{\alpha} \sim \frac{w_{\alpha'}(\epsilon)}{\alpha'}$$

such that $e_{w_\alpha(\epsilon)}(\epsilon) = q_\alpha(\epsilon)$

Case B: Dynamic ϵ

Theorem B Suppose $\epsilon = \epsilon_\alpha \rightarrow 0$ as $\alpha \rightarrow 0$, we have

a) If $F(x)/\{\epsilon H(x)\} \rightarrow 0$ as $\alpha \rightarrow 0$ with x the α quantile of F_ϵ , then

$$q_\alpha(\epsilon) \sim q_{\alpha/\epsilon}(1), \quad ES_\alpha(\epsilon) \sim ES_{\alpha'}(1), \quad \frac{w_\alpha(\epsilon)}{\alpha} \sim \frac{w_{\alpha'}(\epsilon)}{\alpha'}$$

b) If $\epsilon H(x)/F(x) \rightarrow 0$ as $\alpha \rightarrow 0$ with x the α quantile of F_ϵ , then

$$q_\alpha(\epsilon) \sim q_\alpha, \quad ES_\alpha(\epsilon) \sim ES_\alpha, \quad w_\alpha(\epsilon) \sim w_\alpha$$

Case C: Independent Small ϵ

Define the **influence function** (IF) of risk measure ϱ of F_ϵ as

$$IF(\varrho; F, H) = \lim_{\epsilon \rightarrow 0} \frac{\varrho(\epsilon) - \varrho}{\epsilon} = \left. \frac{\partial \varrho(\epsilon)}{\partial \epsilon} \right|_{\epsilon=0}$$

Hence, approximations of $\varrho(\epsilon)$ with infinitesimal contamination level ϵ is given by

$$q_\alpha(\epsilon) \simeq q_\alpha + \epsilon IF(q_\alpha; F, H), \quad ES_\alpha(\epsilon) \simeq ES_\alpha + \epsilon IF(ES_\alpha; F, H)$$

Case C: Influence Function

Theorem C Assume that F has positive continuous differential at its α quantile, and H is continuous at q_α . We have

$$IF(q_\alpha; F, H) = \frac{\alpha - H(q_\alpha)}{F'(q_\alpha)}$$

$$IF(ES_\alpha; F, H) = \frac{q_\alpha \{\alpha - H(q_\alpha)\} + \int_{-\infty}^{q_\alpha} x d\{H(x) - F(x)\}}{\alpha}$$

Example

Example 1 Normal contamination

Consider $F_\epsilon(x) = (1 - \epsilon)\Phi(x) + \epsilon\Phi(x/\sigma)$ with $\sigma > 1$, the scale parameter of the contamination model

- The larger the σ is, the heavier the contamination model is
- For $\epsilon \in (0, 1)$, we have $\varrho_\alpha(\epsilon) \sim \varrho_{\alpha/\epsilon}$
- Let $\epsilon = \alpha^\tau$ with $\tau > 0$ a constant. The larger τ is, the lower the contamination level is, and thus the risk measure might be more robustness

Example

Example 2 Laplace contamination

Consider $F_\epsilon(x) = (1 - \epsilon)\Phi(x) + \epsilon L(x/\sigma)$ with $\sigma > 0$ and L the standard double-sided exponential distribution

$$L(x) = \frac{1}{2} \exp \left\{ -\sqrt{2}|x| \right\}, \quad x \in \mathbb{R}$$

- Essential heavier of Laplace model than normal model
- Approximations based on L might be faster than normal contamination
- In order to obtain non-sensitivity of risk measures, a very smaller $\epsilon = \alpha^\tau$ (and thus a larger τ) is needed

Example

Example 3 Power-like contamination

Consider $F_\epsilon(x) = (1 - \epsilon)\Phi(x) + \epsilon H(x/\sigma)$ with $\sigma > 0$ and H a symmetry distribution with

$$H(x) = \frac{1}{2} \left\{ 1 - \left(1 - \frac{4}{4 + x^2} \right)^{0.5} \right\}, \quad x < 0$$

- Power decaying tail and infinite variance, not useful in practice
- Typical example with expectile coinciding with quantile
- All of the following results are obtained by R codes at

Laplace: $\epsilon = 0.5, \sigma = 1.6$




Figure 2: Approximations based on $N(0,1)$ to its **true values**, indicated by **dotted line** and black line

Laplace: $\alpha = 0.5\%$, $\sigma = 1$




Figure 3: Approximations based on $N(0,1)$ to its **true values**, indicated by **dotted red** and black line

Laplace: $\tau = 1, \sigma = 1.6$




Figure 4: Approximations based on $N(0,1)$ to its **true values**, indicated by **dotted red** and black line

Laplace: $\tau = 0.1, \sigma = 0.95$




Figure 5: Approximations based on $L(\sigma = 0.95)$ to its **true values**, indicated by **dotted line** and black line

RE Based on IF

Approximate of VaR and ES for F_ϵ with small ϵ as

$$\tilde{\varrho}(\epsilon) \stackrel{\text{def}}{=} \varrho + \epsilon IF(\varrho; F, H)$$

Define thus the **relative error** (RE) as

$$\text{RE}(\varrho) = \frac{\tilde{\varrho}(\epsilon) - \varrho(\epsilon)}{\varrho(\epsilon)}$$

◀ Influence function

RE Based on IF

Normal: $\sigma = 2$					
ϵ	0.10	2.10	4.10	6.10	8.10
$RE(q_\alpha)$	-0.02	-0.01	0.01	0.04	0.08
$RE(ES_\alpha)$	0.01	0.01	0.01	0.02	0.03
Laplace: $\sigma = 1.2$					
ϵ	0.10	2.10	4.10	6.10	8.10
$RE(q_\alpha)$	-0.02	-0.01	0.02	0.06	0.11
$RE(ES_\alpha)$	0.01	0.01	0.01	0.02	0.03
Power-like: $\sigma = 1$					
ϵ	0.10	2.10	4.10	6.10	8.10
$RE(q_\alpha)$	-0.02	-0.11	-0.33	-0.71	-1.22
$RE(ES_\alpha)$	0.01	0.03	0.08	0.15	0.26

Table 1: RE of VaR and ES with $\alpha = 0.25$ is in ‰

Tail Analyses via MEF

- Empirical Mean Excess (EME) of $X \sim F$ is given as

$$\hat{m}_X(t) = \frac{\sum_{i=1}^n (X_i - t) \mathbf{I}\{X_i > t\}}{\sum_{i=1}^n \mathbf{I}\{X_i > t\}}, \quad t \text{ large}$$

- Power-like tails as $\bar{F}(x) \sim Cx^{-1/\gamma}$ are implied by

$$\hat{m}_X(t) \sim \frac{\gamma}{1-\gamma} t, \quad \gamma \geq 0$$

- Weibull-like tails as $\bar{F}(x) \sim C \exp\{-x^\tau\}$ are implied by

$$\log \hat{m}_X(t) \sim (1 - \tau) \log t, \quad \tau > 0$$

CRIX: Tail Heaviness Exploration




Figure 6: EME and log EME for daily log returns of CRIX during 20140731–20180101. Lower tail of X is given by the upper tail of $-X$

SRMC—Sensitivity of Risk Measures

Model Choice of CRIX

- Data appears with Laplace tails
- The principal data can be modeled by normal distribution
- Consider the normal-Laplace contamination model with parameter ϵ , $\boldsymbol{\mu} = (\mu_1, \mu_2)$, $\boldsymbol{\sigma} = (\sigma_1, \sigma_2)$

$$F_{\epsilon}(x) = (1 - \epsilon) * \Phi\left(\frac{x - \mu_1}{\sigma_1}\right) + \epsilon * L\left(\frac{x - \mu_2}{\sigma_2}\right), \quad x \in \mathbb{R}$$

Estimation of CRIX

- EM algorithm to estimate parameters involved
- Three periods are considered
 - ▶ A, 20140731–20180101
 - ▶ B, 20140731–20160401
 - ▶ C, 20160402–20180101
- Three methods to estimate VaR and ES
 - ▶ Historical simulations: \widehat{q}_{α}^* , \widehat{ES}_{α}^*
 - ▶ Laplace approximation: $\widehat{q}_{\alpha'}(1)$, $\widehat{ES}_{\alpha'}(1)$
 - ▶ Complete mixture model: $\widehat{q}_{\alpha}(\epsilon)$, $\widehat{ES}_{\alpha}(\epsilon)$

Estimation of CRIX

per.	ϵ	μ_1	μ_2	σ_1	σ_2
A	0.622	0.002	0.004	0.010	0.045
B	0.480	0.001	-0.002	0.014	0.045
C	0.731	0.002	0.008	0.006	0.045

Table 2: Estimated parameters of normal-Laplace contamination

Estimation of CRIX

per.	$\alpha(\%)$	\hat{q}_α^*	$\hat{q}_{\alpha'}(1)$	$\hat{q}_\alpha(\epsilon)$	\widehat{ES}_α^*	$\widehat{ES}_{\alpha'}(1)$	$\widehat{ES}_\alpha(\epsilon)$
A	0.5	0.136	0.127	0.127	0.183	0.125	0.000
	1	0.105	0.105	0.105	0.152	0.138	0.000
	5	0.054	0.055	0.054	0.091	0.085	0.000
B	0.5	0.118	0.125	0.125	0.171	0.184	0.000
	1	0.104	0.103	0.103	0.143	0.130	0.000
	5	0.046	0.052	0.052	0.086	0.094	0.000
C	0.5	0.137	0.128	0.128	0.179	0.128	0.000
	1	0.108	0.106	0.106	0.155	0.130	0.000
	5	0.059	0.055	0.055	0.095	0.080	0.000

Table 3: Estimated VaR, ES based on normal-Laplace model

Conclusions

- Common data-sets such as CRIX follow certain contamination models in Huber's framework
- Practitioners must model carefully the tail feature in risk management
- Theoretical approximations of risk measures are given with illustrated examples

Further work

- How to model the *dynamic* tail features of the data-sets?
- What about the *min-max* problem for these risk measures?
- How about the *statistical estimations* of these models?

How Sensitive are Tail-Related Risk Measures in a Contamination Neighbourhood?

Wolfgang Karl Härdle
Chengxiu Ling



Ladislav von Bortkiewicz
Chair of Statistics
Humboldt-Universität zu Berlin
School of Mathematics & Statistics
Southwest University
crix.berlin swu.edu.cn



References



Brazauskas, V.

Influence functions of empirical nonparametric estimators of net reinsurance premiums

Insurance: Mathematics and Economics 32(1): 115–133, 2003.



Chen, S., Chen, C., Härdle, K. W., Lee, T. Ong, B.

Econometric analysis of a cryptocurrency index for portfolio investment

Handbook of Blockchain, Digital Finance, and Inclusion, Volume 1, Elsevier, pp. 175–206, 2017.

References



Cont, R., Deguest, R. Scandolo, G.

Robustness and sensitivity analysis of risk measurement procedures

Quantitative Finance 10(6): 593–606, 2010.



de Haan, L. Ferreira, A.

Extreme value theory, Springer Series in Operations Research and Financial Engineering

Springer, New York, 2006.

References



Dempster, A., Laird, N. Rubin, D.

Maximum likelihood from incomplete data via the em algorithm

Journal of the Royal Statistical Society. Series B (methodological) pp. 1–38, 1977.



Dierckx, G., Beirlant, J., De Waal, D. Guillou, A.

A new estimation method for Weibull-type tails based on the mean excess function

Journal of Statistical Planning and Inference
139(6): 1905–1920, 2009.

References



Engelke, S. Ivanovs, J.

Robust bounds in multivariate extremes

The Annals of Applied Probability **27**(6): 3706–3734, 2017.



Fermanian, J. Scaillet, O.

Sensitivity analysis of VaR and expected shortfall for portfolios under netting agreements

Journal of Banking & Finance **29**(4): 927–958, 2005.



Ghosh, A.

Divergence based robust estimation of the tail index through an exponential regression model

Statistical Methods & Applications **26**(2): 181–213, 2017.

References



Huber, P.J.

Robust estimation of a location parameter

The Annals of Mathematical Statistics **35**(1): 73–101, 1964.



Kuan, C., Yeh, J. Hsu, Y.

Assessing Value-at-Risk with CARE, the conditional autoregressive expectile models

Journal of Econometrics **150**(2): 261–270, 2009.



McNeil, A., Frey, R. Embrechts, P.

Quantitative Risk Management: Concepts, Techniques and Tools

Princeton university press, 2015.

References

-  Mihoci, A., Härdle, K. W. Chen, C.
TERES-Tail event risk expectile based on shortfall
revised and resubmitted to *Quantitative Finance*, 1–13, 2017.
-  Taylor, J.
Estimating Value-at-Risk and expected shortfall using expectiles
Journal of Financial Econometrics 6(2): 231–252, 2008.
-  Trimborn, S. Härdle, K. W.
CRIX an index for blockchain based currencies
SFB 649 Discussion Paper 2016-021 submitted to *Journal of Empirical Economics*, 2016.

References



Xu, X., Mihoci, A. Härdle, K. W.

ICARE–localizing conditional autoregressive expectiles

Journal of Empirical Economics to appear, 2018.



Zhu, S. Fukushima, M.

Worst-case conditional Value-at-Risk with application to robust portfolio management

Operations Research **57**(5): 1155–1168, 2009.

Expectile and Quantile

- Both are **elicitable**, since e_α and q_α correspond to ϱ_α^2 and ϱ_α^1 , respectively

$$\varrho_\alpha^\gamma = \arg \min_x \alpha E[(X - x)_+]^\gamma + (1 - \alpha) E[(X - x)_-]^\gamma$$

- First-order condition:** setting $x = e_\alpha, q_\alpha$ subsequently

$$\begin{aligned} \alpha E[(X - x)_+] &= (1 - \alpha) E[(X - x)_-] & (2) \\ \alpha \mathbb{P}\{X \geq x\} &= (1 - \alpha) \mathbb{P}\{X \leq x\} \end{aligned}$$

Expectile and Quantile

Specifying $x = e_{w_\alpha}$ in (2), we have

$$\begin{aligned}
 w_\alpha &= \frac{E[(X - x)_-]}{E[(X - x)_-] + E[(X - x)_+]} \\
 &= \frac{E[(X - x)_-]}{2E[(X - x)_-] + E[X] - x}, \quad \text{by using } y_+ = y + y_-
 \end{aligned}$$

with

$$E[(X - x)_-] = xF(x) - \int_{-\infty}^x ydF(y)$$

The desired expression w_α in (1) is thus obtained since $F(x) = \alpha$ due to $e_{w_\alpha} = q_\alpha$ ◀ Common risk measure